103201

232017

103.3.1 Chronic Toxicity

Data Review Number: ES W2

Test: Avian Reproduction Wild Waterfowl

Species: Mallard Duck (Anas platyrhynchos)

Results: Z-9-tricosene was fed to mature Mallard ducks at

dietary concentrations of 0.1 ppm throughout the One-Generation Reproduction Study and had no effect

on the overall reproductive success of birds.

Based on the results of this study, environmental levels of up to 0.01 ppm of Z-9-tricosene do not present a reproductive hazard to the Mallard duck.

Chemical: Muscamone (Technical Z-9-tricosene)

Title: One-Generation Reproduction Study - Mallard Duck

Z-9-Tricosene Final Report

Accession No.: 232017 SAME AS 229393

Study Date: August 28, 1975

Researcher: Wildlife Research Division, Thuron Industries, Inc.

Registrant: Thuron Industries, Inc.

Validation Category: Supplemental

Category Repairability: Yes. The Environmental Safety Section needs to be provided the raw data for penned birds/cage/day/week of eggs set for all listed reproductive parameters. The data is required in order to check the statistical analysis of this data. Test procedures for this study as listed appear adequate.



ADDITIONAL INFORMATION:

Reproductive Data - Mallard Duck	Controls	Z-9-tricos 0.1 (9417%)	
Eggs Laid	909	932	955
Eggs Cracked	15	11	11
Eggs Set*	854	881	904
Viable Embryos	811	832	850
Live Three-Week Embryos	799	818	841
Normal Hatchlings	555	528	591
Fourteen-Day-Old Survivors	553	524	585

^{*} Excludes those cracked and those removed for eggshell thickness analysis. The above differences were not statistically significant (p < 0.05).

	Expected Values	Controls	Z-9-tricoser 0.1 (94.7%) 0.	
Eggs Laid Per Hen In	,			
Eight Weeks	28 - 38	36.4	37.3	38.2
Eggs Cracked of Eggs				
Laid (%)	0.6 - 6%	1.6	1.2	1.2
Viable Embryos of Eggs				
Set (%)	85 - 98%	95	94	94
Live Three-Week Embryos				
of Viable Eggs (%)	97 - 99%	98	98	99
Normal Hatchlings of Live				
Three-Week				
Embryos (%)	50 - 90%	69	64	70
Fourteen-Day-Old Survivors Of Normal	S			
Hatchlings (%)	94 - 99%	99	99	99
Fourteen-Day-Old				
Survivors Per Hen	11 - 14	22.1	21.0	23.4

Statistical analysis based on data in Table la. * The above differences were not statistically significant (p < 0.05).

Eggshell Thickness Data - Mallard Duck Controls $\frac{Z-9-tricosene}{0.1\ (94.7\%)}$ 0.1 (63.1%)

Number of Eggs Analyzed 40 40 40

Mean Shell Thickness (mm) 0.350 0.341 0.346

The above differences were not statistically significant (p < 0.05).

Eggs Land Malland Wo Sign diEFF (POS) M Wildlife INT 6	Z 9 Tricosène Eggo Set. Mallard	Z-9 TricoseNE Viable Embryos
C. 184. control 175. 217. 186. 145.	NO sign diff (P=05) wildlife Int	No sign diff (P=0 wealife Int
181.8 482.56	C. PGM 6	
142. 175. O.lppm 194. 94.7% 208. 213. 4.1	C. 1C. OP 159.59 17 176.	171. 136. 201. Contro 17C. 133.
186.4 666. <i>€</i> 4	155. 204.	162.2 636.56
174. 0.1 ppm. 176. 63.1% AL 198. 232.	17€ 4. 135. 5.	127. Olippin 158. Olippin 174. 94,7%, 187. 198.
191. 512.	170.8 170.8 461.36	166.4 579.44
528165623 (F3.87 2. n.df 12. d.df	461.36	158. 155. Olppm 161. 184. 63.1 % A)
8308. E 55 ² 211.6 T55 ² 8519.6 T6T55 ²	135. 163. O.LPPM 186. 195. 94,7% A.I 204.	17c. 17c. 226.
and the second second	176. 2 652. 56	1268238558 (F3.89
		721C. 152.4 7362.4

; **-**

the second the second of the second s	The state of the s	A CONTRACTOR OF THE PROPERTY O
Z-9 Tricos ene Live Three wish Employed Mallard No. Sign diff (P=0.05	Z.9 Tricosene We Normal Hatching Malland Duch No Sign diff (P:0,04) Whalife Int.	2-9 Tricosene 14 Day OLD Surrows Malland Duck Ne Sign diff (PO.05) Willing Int.
while and	117. 88. 142. control 120. 88.	95. control 94. 75. 95. 55.
135. Control 198. 166. 131.	111. 427.2	68. 47. 69.125 372.859375
159.8 605.36	85. O.L ppm- 111. 94.7% M.L.	25. 68. 72. O.I. ppm
127- 0:1 ppm 151. 0:1 ppm 172. 94.7%A.1 185.	114. 105.6 389.44	87. 94.7%4.1 76. 72. 75. 51.
163.6 480.64	112: Oilppm 95: Oilppm 115: 63:1% A:1.	65, 5 275, 75
157. O.I ppm 158. 634 % A.L 188. 634 % A.L	154. 118. 2	55. 105. 0:1 ppn
185.	396, 56	91. 681% na. 75: 75:
168.2 218.56	. 3952522255 < F 3.89	\$7.000
.1627521924 < F 3,89	6066. 399. 6	73.125 265.609375
6522.8 176.9333334	6465.6	.3341480089 < 4.32 2. 21.
6699.733333		7313.75 232.75
		7546.5

Z-9 TricoseNE Shall Thickness Mallard No Sign diff (PO.05) Wildlife Int.

0. 35 0. 353 **control** 0. 357 0. 351

0.3458 0.00004056

> 0.342 0.316 **0.1 ppm** 0.352 **94.7% A.1** 0.356

0.34C6 0.00020544

> 0.354 0.16 pm 0.356 63.1 % A1 0.374 63.1 % A1 0.346

0.3462 0.00021056

.5649202751 **< F 3.89**

0.0022828 .0002149333 .0024977333